

Client Name \_\_\_\_\_

## Lower Leg Exercise Program

The following routine should be performed \_\_\_\_\_ per day.



This routine can be performed while sitting.

Exercise is important for managing the swelling in your legs.

### Exercise:

- Works your muscles, which assists with moving the lymph fluid away from the swollen area.
- It keeps your joints flexible, maintaining and improving your range of movement.
- It can help you maintain a healthy weight, which can reduce the swelling.

Exercise can also help you feel better about yourself and reduce stress and anxiety.

Make sure you sit up straight with your back well supported.


The following are some simple exercises techniques to get you started. They should **not cause pain**. For further information or if you have any questions talk to your health practitioner.

- 1** Bring your right knee towards your chest.  
Do \_\_\_\_\_ of these.  
Repeat with your left leg.



Practitioner Name: \_\_\_\_\_ Date : \_\_\_\_\_

Client Name \_\_\_\_\_



- 2 With your feet flat on the ground, straighten your right knee and pull your toes towards you.



Hold for \_\_\_\_\_

Do \_\_\_\_\_ of these.

Repeat with your left leg.

- 3 Place your hands just above the crease of your right knee and bend your knee 10 times against your hand.

Repeat with your left leg.



- 7 Slowly pull your right foot up towards your face and then point your foot down. Repeat 10 times.

Repeat with your left foot.

- 8 Slowly turn your right foot in a clockwise direction 10 times. Repeat in an anti-clockwise direction 10 times.

Repeat with your left foot.




Practitioner Name: \_\_\_\_\_ Date: \_\_\_\_\_

Client Name \_\_\_\_\_


- 9 Place your hand just above the crease of your right knee and bend your knee 10 times against your hand.

Repeat with your left leg.




- 10 If your balance is adequate, in the standing position hold onto the kitchen bench and lift your heels up slowly and then return to the ground.

Repeat \_\_\_\_\_



If you can't do this exercise in standing, sit in a chair with your feet flat on the ground. Raise your heels upwards while keeping your toes on the floor. Lower your heels to the ground.

Repeat \_\_\_\_\_



- 11 Try and walk 15 to 20 minutes per day, 3 to 5 days a week. Initially this can be broken up into short walks such as two lots of 10 minutes per day. You might find a pedometer and a log book useful to track your progress. Discuss this with your therapist. Gradually increase your walking each week.

Practitioner Name: \_\_\_\_\_ Date: \_\_\_\_\_